

Structure of Matter

Dr Mohammad Abdur Rashid


Classification of solids

Solids are broadly classified into two types crystalline solids and amorphous solids.
















Simple cubic





Body-centered cubic



Face-centered cubic








Simple cubic

Body-centered cubic

Face-centered cubic



Particle Contributions to Unit Cell



webassign.net

Unit cell

The Simplest Repeating Unit in a Crystal

The simplest repeating unit in a crystal is called a **unit cell**. Each unit cell is defined in terms of **lattice points**, the points in space about which the particles are free to vibrate in a crystal.

Bravais lattice

A Bravais lattice, studied by Auguste Bravais (1850), is an infinite array of discrete points in three dimensional space generated by a set of discrete translation operations described by:


$$\vec{R} = n_1 \vec{a}_1 + n_2 \vec{a}_2 + n_3 \vec{a}_3$$

Where n_i are any integers and \vec{a}_i are known as the primitive vectors which lie in different directions and span the lattice. This discrete set of vectors must be closed under vector addition and subtraction. For any choice of position vector \vec{R} , the lattice looks exactly the same.

$$ec{R} = n_1 ec{a}_1 + n_2 ec{a}_2$$


$$ec{R} = n_1 ec{a}_1 + n_2 ec{a}_2$$
 $1 \quad 0 \quad 0 \quad 1 \quad ec{a}_2 \quad ec{a}_1 \quad ec{a}_2 \quad ec{a}_1 \quad ec{a}_1 \quad ec{a}_2 \quad ec{a}_2 \quad ec{a}_2 \quad ec{a}_1 \quad ec{a}_2 \quad ec{a}_2 \quad ec{a}_2 \quad ec{a}_3 \quad ec{a}_4 \quad ec{a}_4$


Bravais lattice in 3D





epionelynx.wordpress.com


basis + lattice = crystal structure
















The lattice of crystalline quartz (SiO₂)





The Miller indices of a lattice plane are the coordinates of the shortest reciprocal lattice vector normal to the plane, with respect to a specific set of primitive reciprocal lattice vectors. Thus a plane with Miller indices $(h \ k \ l)$, is normal to the reciprocal lattice vector

$$h\vec{b}_1 + k\vec{b}_2 + l\vec{b}_3$$


Miller indices are used to specify directions and planes.

The rules for Miller Indices:

•Determine the intercepts of the face along the crystallographic axes, in terms of unit cell dimensions.

- Take the reciprocals
- Clear fractions
- Reduce to lowest terms

$$(1,2,3) \to (\frac{1}{1}:\frac{1}{2}:\frac{1}{3}) \to (6:3:2) \to (632)$$





Miller Indices are the reciprocals of the parameters of each crystal face.


For Pink Face:
$$\left(\frac{1}{1}, \frac{1}{\infty}, \frac{1}{\infty}\right) \to (100)$$


For Green Face:
$$\left(\frac{1}{\infty}, \frac{1}{\infty}, \frac{1}{1}\right) \to (001)$$


For Yellow Face:
$$\left(\frac{1}{\infty}, \frac{1}{1}, \frac{1}{\infty}\right) \to (010)$$

















Draw your own lattice planes




https://www.doitpoms.ac.uk/tlplib/miller_indices/lattice_draw.php

Band Theory of Solids



Formation of hydrogen molecule





University Physics- Young, Freedman





Energy Bands for Solids







Point defects in a crystal. (a) Vacancy. (b) Interstitial.

(c) Substitutional impurity. (d) Interstitial impurity.


Concepts of Modern Physics – Arthur Beiser







A screw dislocation.

Concepts of Modern Physics - Arthur Beiser







nde-ed.org