

Quantum Mechanics I

PHY 3103

Dr Mohammad Abdur Rashid



Course site

tiny.cc/phy3103

tiny.cc/qm20



References

Quantum Mechanics (6th Ed.) – B.H. Bransden & C.J. Joachain

Quantum Mechanics (2nd Ed.) – Nouredine Zettili



Introduction To Quantum Mechanics (2nd Ed.) – David J. Griffiths




What is light?



Blackbody spectra



Planck Radiation Formula



$$u(\nu) d\nu = \frac{8\pi h}{c^3} \frac{\nu^3 d\nu}{e^{h\nu/kT} - 1}$$



Planck Radiation Formula


$$u(\nu) d\nu = \frac{8\pi h}{c^3} \frac{\nu^3 d\nu}{e^{h\nu/kT} - 1}$$

$$h = 6.626 \times 10^{-34} \text{ J} \cdot \text{s}$$


$$\epsilon_n = nh\nu \quad n = 0, 1, 2, \dots$$



The photoelectric effect





The photoelectric effect




The photoelectric effect

Photoelectric effect

$$h\nu = \text{KE}_{\text{max}} + \phi$$



The photoelectric effect




Work function $\phi = h\nu_0$

What is light?




The Compton effect



$$\lambda' - \lambda = \frac{h}{mc}(1 - \cos \phi)$$

Interference of light



WHAT IS LIGHT?

Both wave and particle



Can particle behave like wave?



De Broglie Waves

A moving body behaves in certain ways as though it has a wave nature

$$\lambda = \frac{h}{p}$$

De Broglie
wavelength

$$\lambda = \frac{h}{\gamma m v}$$

$$\gamma = \frac{1}{\sqrt{1 - v^2/c^2}}$$



Example

Find the de Broglie wavelengths of (a) a 46-g golf ball with a velocity of 30 m/s, and (b) an electron with a velocity of 10^7 m/s.

Solution

(a) Since $v \ll c$, we can let $\gamma = 1$. Hence

$$\lambda = \frac{h}{mv} = \frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s}}{(0.046 \text{ kg})(30 \text{ m/s})} = 4.8 \times 10^{-34} \text{ m}$$

The wavelength of the golf ball is so small compared with its dimensions that we would not expect to find any wave aspects in its behavior.



Example

Find the de Broglie wavelengths of (a) a 46-g golf ball with a velocity of 30 m/s, and (b) an electron with a velocity of 10^7 m/s.

Solution


(b) Again $v \ll c$, so with $m = 9.1 \times 10^{-31}$ kg, we have

$$\lambda = \frac{h}{mv} = \frac{6.63 \times 10^{-34} \text{ J} \cdot \text{s}}{(9.1 \times 10^{-31} \text{ kg})(10^7 \text{ m/s})} = 7.3 \times 10^{-11} \text{ m}$$

The dimensions of atoms are comparable with this figure—the radius of the hydrogen atom, for instance, is 5.3×10^{-11} m. It is therefore not surprising that the wave character of moving electrons is the key to understanding atomic structure and behavior.




Bohr atomic model of a nitrogen atom



© 2011 Encyclopædia Britannica, Inc.



Bohr Atom Model



Condition for
orbital stability

$$mvr = \frac{nh}{2\pi}$$



$$n = 1, 2, 3, \dots$$

Condition for stable orbit


An electron can circle a nucleus only if its orbit contains an integral number of de Broglie wavelengths.

Condition for orbit stability


$$n\lambda = 2\pi r_n \quad n = 1, 2, 3, \dots$$



A fractional number of wavelengths cannot persist because destructive interference will occur



Energy levels of hydrogen atom




Jim Al-Khalili


- Quantum Life: How Physics Can Revolutionise Biology
- Double Slit Experiment explained! by Jim Al-Khalili


































Ri





Are you shocked?



Those who are not shocked when they first come across quantum mechanics cannot possibly have understood it.

Niels Bohr

